



# Nuclear Astrophysics

## I. Stellar burning

Karlheinz Langanke


GSI & TU Darmstadt

Aarhus, October 6-10, 2008

# What is nuclear astrophysics?

Nuclear astrophysics aims at understanding the nuclear processes that take place in the universe. These nuclear processes generate energy in stars and contribute to the nucleosynthesis of the elements.


## 3. The solar abundance distribution



### solar abundances:


Elemental (and isotopic) composition of Galaxy at location of solar system at the time of its formation

Hydrogen mass fraction	$X = 0.71$
Helium mass fraction	$Y = 0.28$
Metallicity (mass fraction of everything else)	$Z = 0.019$
Heavy Elements (beyond Nickel) mass fraction	$4E-6$




N. Grevesse and A. J. Sauval, *Space Science Reviews* **85**, 161

# Hoyle's cosmic cycle



# Nucleosynthesis processes

In 1957: Burbidge, Burbidge, Fowler, Hoyle, [Rev. Mod. Phys. **29**, 547 (1957)] suggested the synthesis of the elements in stars.




# Where does the energy come from?

Energy comes from nuclear reactions in the core.



$$E = mc^2$$



The Sun converts 600 million tons of hydrogen into 596 million tons of helium every second. The difference in mass is converted into energy. The Sun will continue burning hydrogen during 5 billions years. Energy released by H-burning:

$$6.45 \times 10^{18} \text{ erg g}^{-1}$$

$$\text{Solar Luminosity: } 3.846 \times 10^{33} \text{ erg s}^{-1}$$

# Types of processes

**Transfer** (strong interaction)

$$^{15}\text{N}(p, \alpha)^{12}\text{C}, \quad \sigma \simeq 0.5 \text{ b at } E = 2.0 \text{ MeV}$$

**Capture** (electromagnetic interaction)

$$^3\text{He}(\alpha, \gamma)^7\text{Be}, \quad \sigma \simeq 10^{-6} \text{ b at } E = 2.0 \text{ MeV}$$

**Weak** (weak interaction)

$$p(p, e^+ \nu)d, \quad \sigma \simeq 10^{-20} \text{ b at } E = 2.0 \text{ MeV}$$

# Stellar reaction rate

Consider  $N_a$  and  $N_b$  particles per cubic centimeter of particle types  $a$  and  $b$ . The rate of nuclear reactions is given by:

$$r = N_a N_b \sigma(v) v$$

In stellar environment the velocity (energy) of particles follows a thermal distribution that depends on the type of particles.

- Nuclei (Maxwell-Boltzmann):  $\phi(v) = N 4\pi v^2 \left(\frac{m}{2\pi kT}\right)^{3/2} \exp\left(-\frac{mv^2}{2kT}\right)$

The product  $\sigma v$  has to be averaged over the velocity distribution  $\phi(v)$


$$\langle \sigma v \rangle = \int_0^\infty \int_0^\infty \phi(v_a) \phi(v_b) \sigma(v) v dv_a dv_b$$

Changing to center-of-mass coordinates, integrating over the cm-velocity and using  $E = \mu v^2/2$

$$\langle \sigma v \rangle = \left(\frac{8}{\pi\mu}\right)^{1/2} \frac{1}{(kT)^{3/2}} \int_0^\infty \sigma(E) E \exp\left(-\frac{E}{kT}\right) dE$$

# Charged-particle cross section



Stars' interior is a plasma made of charged particles (nuclei, electron). Nuclear reactions proceed by tunnel effect. For  $p + p$  reaction Coulomb barrier 550 keV, but the typical energy in the sun is only 1.35 keV.



$$\text{cross section: } \sigma(E) = \frac{1}{E} S(E) e^{-2\pi\eta}; \quad \eta = \frac{Z_1 Z_2 e^2}{\hbar} \sqrt{\frac{\mu}{2E}} = \frac{b}{E^{1/2}}$$

# Astrophysical S factor


S factor allows accurate extrapolations to low energy.



# Gamow window

Using definition of S factor:

$$\langle \sigma v \rangle = \left( \frac{8}{\pi \mu} \right)^{1/2} \frac{1}{(kT)^{3/2}} \int_0^{\infty} S(E) \exp \left[ -\frac{E}{kT} - \frac{b}{E^{1/2}} \right] dE$$



# Gamow window

Assuming that S factor is constant over the Gamow window and approximating the integrand by a Gaussian one gets:

$$\langle \sigma v \rangle = \left( \frac{2}{\mu} \right)^{1/2} \frac{\Delta}{(kT)^{3/2}} S(E_0) \exp \left( -\frac{3E_0}{kT} \right)$$

$$E_0 = 1.22[\text{keV}](Z_1^2 Z_2^2 \mu T_6^2)^{1/3}$$

$$\Delta = 0.749[\text{keV}](Z_1^2 Z_2^2 \mu T_6^5)^{1/6}$$

( $T_x$  measures the temperature in  $10^x$  K.)


Examples for solar conditions:

reaction	$E_0$ [keV]	$\Delta/2$ [keV]	$I_{\text{max}}$	T dependence of $\langle \sigma v \rangle$
p+p	5.9	3.2	$1.1 \times 10^{-6}$	$T^{3.9}$
p+ $^{14}\text{N}$	26.5	6.8	$1.8 \times 10^{-27}$	$T^{20}$
$\alpha$ + $^{12}\text{C}$	56.0	9.8	$3.0 \times 10^{-57}$	$T^{42}$
$^{16}\text{O}$ + $^{16}\text{O}$	237.0	20.2	$6.2 \times 10^{-239}$	$T^{182}$

It depends very sensitively on temperature!

# The p-p chain


Step 1:  $p + p \rightarrow {}^2\text{He}$  (not possible)



Step 2:  $d + p \rightarrow {}^3\text{He}$



Step 3:  ${}^3\text{He} + p \rightarrow {}^4\text{Li}$  ( ${}^4\text{Li}$  is unbound)



$d + d$  is not going, because  $Y_d$  is extremely small and  $d + p$  leads to rapid destruction.

${}^3\text{He} + {}^3\text{He}$  works, because  $Y_{{}^3\text{He}}$  increases as nothing destroys it.

# The relevant S-factors

$p(p, e^+ \nu_e)d$ :  $S_{11}(0) = (4.00 \pm 0.05) \times 10^{-25} \text{ MeVb}$

calculated

$p(d, \gamma)^3\text{He}$ :  $S_{12}(0) = 2.5 \times 10^{-7} \text{ MeVb}$

measured at LUNA

$^3\text{He}(^3\text{He}, 2p)^4\text{He}$ :  $S_{33}(0) = 5.4 \text{ MeVb}$

measured at LUNA



Laboratory Underground for Nuclear Astrophysics (Gran Sasso) nframe

# Burning of Deuterium



Deuterons are burnt by the reaction  $d(p, \gamma)^3\text{He}$ :

$$\begin{aligned}\frac{dD}{dt} &= r_{11} - r_{12} \\ &= \frac{H^2}{2} \langle \sigma v \rangle_{11} - HD \langle \sigma v \rangle_{12}\end{aligned}$$

In equilibrium ( $\frac{dD}{dt} = 0$ ) one has

$$\begin{aligned}\left(\frac{D}{H}\right)_e &= \frac{\langle \sigma v \rangle_{11}}{2 \langle \sigma v \rangle_{12}} \\ (D/H)_e &= 5.6 \times 10^{-18} \text{ for } T_6 = 5\end{aligned}$$

# The ppl chain




# $^4\text{He}$ as catalyst

$^4\text{He}$  can act as catalyst initializing the ppII and ppIII chains.  
With which nucleus will  $^4\text{He}$  fuse?

- protons:  
the fusion of  $^4\text{He}$  and protons lead to  $^5\text{Li}$  which is unbound.
- deuterons:  
the fusion of deuterons with  $^4\text{He}$  can make stable  $^6\text{Li}$ ; however, the deuteron abundance is too low for this reaction to be significant
- $^3\text{He}$ :  
 $^3\text{He}$  and  $^4\text{He}$  can fuse to  $^7\text{Be}$ . This is indeed the break-out reaction from the ppI chain.

Once  $^7\text{Be}$  is produced, it can either decay by electron capture or fuse with a proton. Thus, the reaction sequence branches at  $^7\text{Be}$  into the ppII and ppIII chains.

# The solar pp chains





# Hydrogen burning: pp-chains vs CNO cycle

Slowest reaction determines efficiency (energy production) of chain:

## pp-chains:

p+p fusion, mediated by weak interaction

## CNO cycle:

$^{14}\text{N}+\text{p}$ , largest Coulomb barrier, mediated by electromagnetic interaction (in contrast to strong interaction in  $^{15}\text{N}+\text{p}$ )


Temperature dependence quite different:

$$\langle \sigma v \rangle \sim T^{(\tau-2)/3}$$

$$\text{with } \tau = \frac{3E_0}{kT}; E_0 = 1.22[\text{keV}](Z_1^2 Z_2^2 \mu T_6^2)^{1/3}$$

$$\text{At } T_6 = 15 \text{ (solar core): } \langle \sigma v \rangle \sim T^{3.9} \text{ (pp); } \langle \sigma v \rangle \sim T^{20} \text{ (CNO)}$$

# Energy generation: CNO cycle vs pp-chains



# Consequences

- stars slightly heavier than the Sun burn hydrogen via CNO cycle
- this goes significantly faster; such stars have much shorter lifetimes

mass [ $M_{\odot}$ ]	timescale [y]
0.4	$2 \times 10^{11}$
0.8	$1.4 \times 10^{10}$
1.0	$1 \times 10^{10}$
1.1	$9 \times 10^9$
1.7	$2.7 \times 10^9$
3.0	$2.2 \times 10^8$
5.0	$6 \times 10^7$
9.0	$2 \times 10^7$
16.0	$1 \times 10^7$
25.0	$7 \times 10^6$
40.0	$1 \times 10^6$

hydrogen burning timescales depend strongly on mass. Stars slightly heavier than the Sun burn hydrogen by CNO cycle.

# End of hydrogen core burning

When the hydrogen fuel in the core gets exhausted, an isothermal core of about 8% of the stellar mass can develop in the center. Continuous hydrogen burning adds to the core mass which eventually rises over the Schönberg-Chandrasekhar mass limit. Then the core's temperature (and density) rise. Finally the central temperature is high enough ( $T_c \approx 10^8$  K) to ignite [helium core burning](#).

Hydrogen burning continues in a shell outside the helium core. This ([hydrogen shell burning](#)) occurs at higher temperatures than hydrogen core burning.

# Which reaction can start helium burning?


Consider a supply of protons and  ${}^4\text{He}$ .

We first note again that  ${}^5\text{Li}$  is unbound. Although this nucleus is continuously formed by  $p+{}^4\text{He}$  reactions, the scattering is elastic and the formed  ${}^5\text{Li}$  nuclei decay within  $10^{-22}$  s.

As a consequence  ${}^4\text{He}$  'survives' in the core until sufficiently large temperatures are achieved to overcome the larger Coulomb barrier between  ${}^4\text{He}$  nuclei. Unfortunately the  ${}^8\text{Be}$  ground state, formed by elastic  ${}^4\text{He}+{}^4\text{He}$  scattering, is a resonance too and decays within  $10^{-16}$  s back to two  ${}^4\text{He}$  nuclei.


# The Salpeter-Hoyle suggestion

In 1952 Salpeter pointed out that the  ${}^8\text{Be}$  lifetime might be sufficiently large that there is a chance that it captures another  ${}^4\text{He}$  nucleus. This *triple-alpha reaction*




can then form  ${}^{12}\text{C}$  and supply energy. However, the simultaneous collision of 3  ${}^4\text{He}$  ( $\alpha$ -particles) is too rare to give the burning rate necessary in stellar models. So Hoyle predicted a resonance in  ${}^{12}\text{C}$  to speed up the collision. And indeed, this *Hoyle state* was experimentally observed shortly after its prediction.

${}^{12}\text{C}$  can then react with another  ${}^4\text{He}$  nucleus forming  ${}^{16}\text{O}$  via




These two reactions make up helium burning.

## Critical Reactions in He-burning



Energy source in stellar He burning  
Energy release determined by associated reaction rates

# Influence of $\alpha+^{12}\text{C}$ on nucleosynthesis



# At the end of helium burning

Nucleosynthesis yields from stars may be divided into production by stars above or below  $9M_{\odot}$ .

- **stars with  $M \lesssim 9M_{\odot}$**   
the stars are expected to shed their envelopes during helium burning and become white dwarfs. Most of the matter returned to the ISM is unprocessed.
- **stars with  $M > 9M_{\odot}$**   
these stars will ignite carbon burning under non-degenerate conditions. The subsequent evolution proceeds in most cases to core collapse. These stars make the bulk of newly processed matter that is returned to the ISM.

# Carbon Burning



## Burning conditions:

for stars  $> 8 M_{\odot}$  (solar masses) (ZAMS)

$T \sim 600\text{-}700 \text{ Mio}$

$\rho \sim 10^5\text{-}10^6 \text{ g/cm}^3$

## Major reaction sequences:



dominates  
by far

of course p's, n's, and a's are recaptured ...  $^{23}\text{Mg}$  can b-decay into  $^{23}\text{Na}$

## Composition at the end of burning:

mainly  $^{20}\text{Ne}$ ,  $^{24}\text{Mg}$ , with some  $^{21,22}\text{Ne}$ ,  $^{23}\text{Na}$ ,  $^{24,25,26}\text{Mg}$ ,  $^{26,27}\text{Al}$

of course  $^{16}\text{O}$  is still present in quantities comparable with  $^{20}\text{Ne}$  (not burning ... yet) <sub>21</sub>

# Neon Burning

Neon burning is very similar to carbon burning.

## Burning conditions:

for stars  $> 12 M_{\odot}$  (solar masses) (ZAMS)

$T \sim 1.3\text{-}1.7 \text{ Bio K}$

$\rho \sim 10^6 \text{ g/cm}^3$

## Why would neon burn before oxygen ???

Answer:


Temperatures are sufficiently high to initiate **photodisintegration** of  $^{20}\text{Ne}$



this is followed by (using the liberated helium)



so net effect:




22

# Oxygen Burning

## Burning conditions:

$$T \sim 2 \text{ Bio}$$
$$\rho \sim 10^7 \text{ g/cm}^3$$

## Major reaction sequences:



plus recapture of n,p,d, $\alpha$

## Main products:

**${}^{28}\text{Si}$ ,  ${}^{32}\text{S}$  (90%)** and some  ${}^{33,34}\text{S}$ ,  ${}^{35,37}\text{Cl}$ ,  ${}^{36,38}\text{Ar}$ ,  ${}^{39,41}\text{K}$ ,  ${}^{40,42}\text{Ca}$

# Silicon Burning

Silicon burning is very similar to oxygen burning.

## Burning conditions:

$T \sim 3-4 \text{ Bio}$

$\rho \sim 10^9 \text{ g/cm}^3$

## Reaction sequences:

- Silicon burning is fundamentally different to all other burning stages.
- **Complex network of fast  $(\gamma, n)$ ,  $(\gamma, p)$ ,  $(\gamma, \alpha)$ ,  $(n, \gamma)$ ,  $(p, \gamma)$ , and  $(\alpha, \gamma)$  reactions**
- The net effect of Si burning is:  $2 \text{ }^{28}\text{Si} \rightarrow \text{}^{56}\text{Ni}$ ,

## need new concept to describe burning:

**Nuclear Statistical Equilibrium (NSE)**

**Quasi Statistical Equilibrium (QSE)**

## Nuclear burning stages

(e.g., 20 solar mass star)

Fuel	Main Product	Secondary Product	T (10 <sup>9</sup> K)	Time (yr)	Main Reaction
H	He	<sup>14</sup> N	0.02	10 <sup>7</sup>	<sup>CNO</sup> 4 H → <sup>4</sup> He
He	O, C	<sup>18</sup> O, <sup>22</sup> Ne s-process	0.2	10 <sup>6</sup>	3 He <sup>4</sup> → <sup>12</sup> C <sup>12</sup> C(α,γ) <sup>16</sup> O
C	Ne, Mg	Na	0.8	10 <sup>3</sup>	<sup>12</sup> C + <sup>12</sup> C
Ne	O, Mg	Al, P	1.5	3	<sup>20</sup> Ne(γ,α) <sup>16</sup> O <sup>20</sup> Ne(α,γ) <sup>24</sup> Mg
O	Si, S	Cl, Ar, K, Ca	2.0	0.8	<sup>16</sup> O + <sup>16</sup> O
Si	Fe	Ti, V, Cr, Mn, Co, Ni	3.5	0.02	<sup>28</sup> Si(γ,α)...

# Kippenhahn diagram for a $22 M_{\odot}$ star

(A. Heger and S. Woosley)

