

Motion in the Schwarzschild metric

Motion in the Schwarzschild metric reveals several of the unusual consequences of general relativity:

1. Utmost *relativity of measurements*: it takes finite proper time for a body to fall onto the Schwarzschild radius, yet for the outside observer it takes infinite time;
2. There exist *gravitational singularities* (geodesic incompleteness) in general relativity: some trajectories cannot be extended beyond a certain point. Gravitational singularities—unlike coordinate singularities—do not depend on the coordinate system and cannot be removed by coordinate transformation. The gravitational field becomes infinitely large at a gravitational singularity.
3. There exist *event horizons* in general relativity – the hyper-surfaces in time-space which can only be crossed in one direction.

Lemaitre coordinates

In the Schwarzschild metric around a body with the gravitational radius r_g ,

$$ds^2 = \left(1 - \frac{r_g}{r}\right) dt^2 - \frac{dr^2}{1 - \frac{r_g}{r}} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (1)$$

there is a singularity at the gravitational radius, $r = r_g$. Under the gravitational radius the coordinate r becomes time-like and t becomes space-like.

However, it turns out to be not a physical singularity, but rather an artifact of the (incorrect) assumption that a static Schwarzschild coordinates can be realized under the gravitational radius with material bodies. Such removable singularities are called *coordinate singularities*.

A transformation to the Lemaitre coordinates τ, ρ

$$d\tau = dt + \sqrt{\frac{r_g}{r}} \frac{dr}{1 - \frac{r_g}{r}} , \quad (2)$$

$$d\rho = dt + \sqrt{\frac{r}{r_g}} \frac{dr}{1 - \frac{r_g}{r}} , \quad (3)$$

leads to the Lemaitre metric, where the singularity at r_g is removed¹,

$$ds^2 = d\tau^2 - \frac{r_g}{r} d\rho^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (4)$$

where $r = [\frac{3}{2}(\rho - \tau)]^{2/3} r_g^{1/3}$. The latter is obtained by integrating

$$d\rho - d\tau = \sqrt{\frac{r}{r_g}} dr , \quad (5)$$

which is the difference between (3) and (2).

The Lemaitre coordinates are synchronous² and are thus realized by a system of clocks in a free radial fall from infinity towards the origin.

Radial fall towards the origin

For a free falling body $d\rho = 0$ and equation (3) gives

$$dt = -\sqrt{\frac{r}{r_g}} \frac{1}{\left(1 - \frac{r_g}{r}\right)} dr . \quad (6)$$

¹ there remains a genuine singularity at the origin.

² the metric has the form $ds^2 = d\tau^2 + g_{\alpha\beta} dx^\alpha dx^\beta$.

Approaching the Schwarzschild radius, in the region $r \gtrsim r_g$, we have in the lowest order in $(r - r_g)/r_g$,

$$dt = -\frac{r_g}{r - r_g} dr, \Rightarrow \frac{r - r_g}{r_1 - r_g} = e^{-\frac{t - t_1}{r_g}}. \quad (7)$$

Apparently, it takes a free falling body infinitely long t -time—the time used by the outer observer—to reach the Schwarzschild radius.

On the contrary, a free falling Lemaitre clock moves from some radius r_1 to a smaller radius r_2 —which can well be the gravitational radius or even the origin—within finite τ -time $\Delta\tau_{12}$. Indeed, setting $d\rho = 0$ in (5) gives

$$\Delta\tau_{12} = -\int_{r_1}^{r_2} \sqrt{\frac{r}{r_g}} dr = \frac{2}{3} \left(\frac{r_1^{3/2} - r_2^{3/2}}{r_g^{1/2}} \right). \quad (8)$$

Event horizons and black holes

Along the trajectory of a radial light ray

$$ds^2 = d\tau^2 - \frac{r_g}{r} d\rho^2 = 0, \quad (9)$$

which gives

$$d\rho = \pm \sqrt{\frac{r}{r_g}} d\tau, \quad (10)$$

where plus and minus describe the rays of light sent correspondingly up and down.

Isolating $d\rho$ in (5) and inserting the result into (10) shows that along the trajectory

$$dr = \left(\pm 1 - \sqrt{\frac{r_g}{r}} \right) d\tau. \quad (11)$$

Apparently if $r < r_g$ then there is always $dr < 0$ and thus the light rays emitted radially inwards and outwards both end up at the origin. In other words no signal can escape from inside the gravitational radius. This phenomenon is called the event horizon.

Therefore a massive object with a size less than the gravitational radius, called a black hole, is completely under the event horizon and its interior is totally invisible.

The trajectories of massive bodies and light rays inside the gravitational radius both end up in the origin where they cannot be extended any further.

The black holes can possibly be detected through their interaction with the matter outside the event horizon.

Exercises

1. In the Schwarzschild metric a body is falling free radially toward the center. What is its coordinate velocity dr/dt at radius r ? What is its locally measured velocity at the same place? Hints: in the Schwarzschild metric the locally measured radial length is given as $d\hat{r}^2 = (1 - \frac{2M}{r})^{-1} dr^2$ and the locally measured time is given as $d\hat{t}^2 = (1 - \frac{2M}{r}) dt^2$.
2. A radio transmitter is falling free radially toward a black hole. When the transmitter is approaching the gravitational radius an outside observer measures its radio signal to be red-shifted as $\exp(-\lambda t)$. Estimate the mass of the black hole from the measured λ . Hints: $\omega_0 = \omega/\sqrt{g_{00}}$, $r - R = (r_1 - R) \exp(-(t - t_1)/R)$.
3. Calculate the proper time it takes for a Lemaitre clock to fall from the gravitational radius to the center of a black hole. For a black hole with the solar mass specify this time in seconds.

4. Show that in Newtonian mechanics the circular planetary orbits around stars are stable against small radial perturbations. Show that in general relativity circular orbits are stable only if $r > 6M$. Hints: consider a circular orbit with a small perturbation, $u = u_0 + \delta u$; derive the equation for δu in the lowest order; investigate whether the perturbation remains small or increases.
5. Show that equatorial orbits in the Schwarzschild metric are stable. Hint: consider an equatorial orbit with a small perturbation, $\theta = \pi/2 + \delta\theta$; derive the equation for $\delta\theta$ in the lowest order; show that the perturbation remains small.