

## Classical tests of general relativity

Einstein proposed three tests of general relativity in 1916 (subsequently called the classical tests):

- The anomalous advance of Mercury's perihelion;
- The deflection of light by the sun;
- The gravitational red-shift of light.

Of these tests, only the perihelion advance of Mercury was known prior to Einstein's final publication of general relativity in 1916. The subsequent experimental confirmation of his other predictions, especially the first measurements of the deflection of light by the sun in 1919, firmly established general relativity as a mainstream theory.

### Mercury perihelion advance

In the 19th century it was discovered that interplanetary perturbations cannot account fully for the turning rate of the Mercury's orbit. About 43 arc-seconds per century remained unexplained. The general theory of relativity exactly accounts for this discrepancy.

The Newtonian equation for the equatorial motion of a planet around a star with mass  $M$  is given as

$$u'' + u = \frac{M}{J^2}, \quad (1)$$

where  $u = 1/r$ ,  $u' = du/d\varphi$ ,  $\varphi$  is the azimuth angle, and  $J$  is a constant. This equation has a periodic elliptic solution with the angular period  $2\pi$ ,

$$u = A \cos(\varphi - \varphi_0) + \frac{M}{J^2}, \quad (2)$$

where  $A$  and  $\varphi_0$  are constants.

The corresponding relativistic equation,

$$u'' + u = \frac{M}{J^2} + 3Mu^2, \quad (3)$$

has an additional relativistic term  $3Mu^2$  which causes the perihelion to shift as illustrated on Figure 1.




Figure 1: Apselion shift of a planet orbiting a star.

The small correction,  $\epsilon \ll 1$ , to the angular frequency can be found by searching for a solution in the form

$$u = A \cos[(1 + \epsilon)\varphi] + B, \quad (4)$$

where  $A$  and  $B$  are constants. Setting this into equation (3) and collecting lowest order terms with powers of  $\cos[(1 + \epsilon)\varphi]$  gives

$$-A2\epsilon \cos[(1 + \epsilon)\varphi] = 3M2AB \cos[(1 + \epsilon)\varphi]. \quad (5)$$

The constant  $B$  has to be taken here in the lowest order,  $B = M/J^2$ , which gives

$$\epsilon = -\frac{3M^2}{J^2}. \quad (6)$$

The angular distance between two perihelia,  $\Delta\varphi$ , is determined by the equation  $(1+\epsilon)\Delta\varphi = 2\pi$ , which gives (in lowest order in  $\epsilon$ )  $\Delta\varphi = 2\pi - 2\pi\epsilon$ . Correspondingly the shift of the orbit,  $\delta\varphi$ , is given as

$$\delta\varphi = -2\pi\epsilon = 2\pi\frac{3M^2}{J^2}. \quad (7)$$

This accounts precisely for the unexplained advance of the Mercury's orbit.

## Bending of light

General relativity predicts apparent bending of light rays passing through gravitational fields. The bending was first observed in 1919 by A.S. Eddington during a total eclipse when stellar images near the occulted disk of the Sun appeared displaced by some arc-seconds from their usual locations in the sky. Later more precise experiments have unambiguously shown that the amount of deflection agrees with the prediction of general relativity.

The *Einstein ring* is an example of the deflection of light from distant galaxies by nearby objects.

In the Newtonian theory the light rays travel along straight lines described by the equation  $u'' + u = 0$  with the (straight-line) solution  $u = A\cos(\varphi - \varphi_0)$ . The corresponding relativistic equation,

$$u'' + u = 3Mu^2, \quad (8)$$

has an additional term,  $3Mu^2$ , which causes the trajectory of light to deflect from the straight line. Assuming the solution in the form  $u = A\cos\varphi + \epsilon(\varphi)$ , where  $\epsilon(\varphi)$  is a small correction, gives

$$\epsilon'' + \epsilon = 3MA^2\cos^2\varphi. \quad (9)$$

Assuming  $\epsilon(\varphi) = C\cos^2\varphi + D$ , where  $C$  and  $D$  are constants, gives

$$\epsilon(\varphi) = MA^2(2 - \cos^2\varphi). \quad (10)$$

The incoming and outgoing rays, where  $r = \infty$  and  $u = 0$ , correspond to the angles  $\varphi_\infty$  which are the solutions to the equation  $u(\varphi_\infty) = 0$ . Searching for the solution perturbatively in the form  $\varphi_\infty = \pi/2 + \delta\varphi$  gives  $\delta\varphi = 2MA$ .

Thus, the relativistic deflection angle  $\Delta\varphi$  between the in-going and out-going rays is

$$\Delta\varphi = 2\delta\varphi = 4MA = \frac{4M}{r_0} = 2\frac{r_g}{r_0}, \quad (11)$$

where  $r_0$  is the closest distance between the ray and the central body.

## Gravitational red-shift

Gravitational red shift is a change of the frequency of the electro-magnetic radiation as it passes through a gravitational field. It is a direct consequence of the equivalence principle.

The connection between the proper time interval  $\Delta\tau$  and the world time interval  $\Delta t$  (here we only consider stationary gravitational fields where such world time can be introduced) is

$$\Delta\tau = \sqrt{g_{00}}\Delta t. \quad (12)$$

Since frequencies are inversely proportional to the time intervals the corresponding connection between world frequency  $\omega_0$  and the locally measured frequency  $\omega$  is

$$\omega = \frac{\omega_0}{\sqrt{g_{00}}}. \quad (13)$$

In a weak gravitational field  $g_{00} = 1 + 2\phi/c^2$  and therefore  $\omega = \omega_0(1 - \phi/c^2)$ . A photon emitted from a point with  $\phi_1$  and received at a point with  $\phi_2$  will be shifted by

$$\Delta\omega = \frac{\phi_1 - \phi_2}{c^2}\omega. \quad (14)$$

The famous experiment that verified the gravitational red-shift is generally called the Pound-Rebka-Snider experiment where the Mössbauer effect was used to accurately measure the change of frequency of a photon travelling upwards 22 m in the Earth's gravitational field.

## Exercises

1. Derive the Kepler's law (the relation between the orbit's period and the radius) for a circular orbit in Schwarzschild metric.

Answer: like in Newtonian theory,  $\omega^2 = M/r^3$ .

Hint: the period is equal  $2\pi/\omega$ , where  $\omega = d\varphi/dt$  is the angular frequency which can be found from the geodesics  $Du^r = 0$ .

2. Consider a nearly circular orbit of a planet around a star in Newtonian mechanics and in General Relativity. Derive the equation for a small radial perturbation of the orbit and find its angular period. Relate to the post-Newtonian perihelion precession.
3. In Newtonian mechanics consider a planet rotating around a star which is slightly non-spherical (a bit of a pancake-like due to the centrifugal forces of the star's rotation), such that the classical Newtonian gravitational potential in the equatorial plane is given as (explain the absence of the  $1/r^2$  term)

$$\phi(r) = -\frac{M}{r} + \frac{MJ_2R^2}{r^3}, \quad (15)$$

where  $J_2$  is the quadrupole moment of the star (presumed very small), and  $R$  is the (equatorial) radius of the star. In Newtonian mechanics calculate the precession of the perihelion of the orbit of the planet. For simplicity you may assume that the orbit is nearly circular.

The quadrupole moment of our sun is estimated<sup>1</sup> to be  $J_2^\odot \approx 2 \times 10^{-7}$ . Can this explain the anomalous precession of Mercury's orbit?

---

<sup>1</sup>Stephany Godier and Jean-Pierre Rozelot, *Quadrupole moment of the Sun: Gravitational and rotational potentials*, Astron. Astrophys. 350, 310–317 (1999); S. Pireaux, J.P. Rozelot, S. Godier, *Solar quadrupole moment and purely relativistic gravitation contributions to Mercury's perihelion Advance*, Astrophys.Space Sci. 284 (2003) 1159–1194 (arXiv:astro-ph/0109032).