



Petar Stipanović





Ground state properties of weakly bound few-body systems

- collaborators –



Leandra Vranješ Markić

Jordi Boronat



EFB23, Aarhus, Denmark, 9th of August 2016.


>> Outline

- researching clusters of few particles

- dimers



- trimers



- tetramers




- ...

- halo state

- energy-size scaling


- universalities

- structural properties




>> Intro: Halo states

- quantum halo states – quantum effects very significant



>> Intro: Halo states



- first were noticed in nuclear physics - dimer



Credit: Ken-ichiro Yoneda, RIKEN Nishina Center for Accelerator-Based Science

>> Intro: Halo states

- characteristics: weakly bound & wide systems



>> Intro: Dimers

? is there unique relationship between physical properties of halo dimers

- dimers - 16 orders of magnit.



- scaled quantities (R ?)

- $x_E = \mu B R^2 / \hbar^2$

- $y_r = \langle r^2 \rangle / R^2$

- halo states

- $y_r = \langle r^2 \rangle / R^2 \geq 2$





>> Intro: Trimers

- trimer properties were scaled:

- $X_E = mB\rho_0^2/\hbar^2$
- $Y_r = \langle \rho^2 \rangle / \rho_0^2$

$$\mu r^2 \downarrow$$
$$m\rho^2 = \frac{1}{M} \sum_{i < k} m_i m_k r_{ik}^2$$


$$M = m_1 + m_2 + \mathbf{m}_3$$




>> Intro: Trimers

- larger number of atomic quantum clusters is missing

? are different-type 3mer
properties related




Phys. Scr. 2013, 014001 (2013) [Riisager]




>> Intro: Trimers

- larger number of atomic quantum clusters is missing

? are different-type 3mer properties related




Phys. Scr. 2013, 014001 (2013) [Riisager]




>> Intro: Trimers

- larger number of atomic quantum clusters is missing

? are different-type 3mer
properties related




Phys. Scr. 2013, 014001 (2013) [Riisager]




>> Intro: Trimers

- larger number of atomic quantum clusters is missing

? are different-type 3mer
properties related




Phys. Scr. 2013, 014001 (2013) [Riisager]



>> Intro: Trimers



- larger number of atomic quantum clusters is missing
- ? are different-type 3mer properties related
- objective: clarify ratio of scaled size and scaled energy
- very important
 - ✗ accuracy of V
 - ✓ precise $E, \langle r_{ij}^2 \rangle$
- method
 - ✓ DMC

Phys. Scr. 2013, 014001 (2013) [Riisager]



>> METHOD: Scaling

- e.g. 2 definitions of R for scaling: $\langle r^2 \rangle / R^2$; $\mu B R^2 / \hbar^2$
 - classical turning point



>> Dimer scaling

$$y_r = r^2 R^{-2}$$

10²

10

1

y_r⁸

$$R = R_v$$

10⁻²

10⁻¹

1

10

$$x_E = \mu B R^2 \hbar^{-2}$$

× ⁴He₂

[a] Rev. Mod. Phys. 76, 215 (2004) [Jensen et al.]

[a] Phys. Scr. 2013, 014001 (2013) [Riisager]

³H ×

d ×
¹¹Be *

³¹Ne

¹⁹C

T [a] ✕
E [a] ●

¹⁵C

³⁵Mg

$$R = R_e$$


10⁻²


10⁻¹


1


10


$$x_E = \mu B R^2 \hbar^{-2}$$






















































































































>> Structure - DMC - pure estimators


- different distribution functions





>> Structure - experiments



>> Structure - experiments



Nature Communications 5, 5765 (2014)




Maksim Kunitski *et al.*, Science 348, 551 (2015)

>> Structure - DMC(V) vs experiment

[a] J. Voigtsberger *et al.*, Nature Comm. 5, 5765 (2014)

[b] Maksim Kunitski *et al.*, Science 348, 551 (2015)



${}^4\text{He}_3$



>> Structure - DMC(V) vs experiment



[a] J. Voigtsberger *et al.*, Nature Comm. 5, 5765 (2014)

[b] Maksim Kunitski *et al.*, Science 348, 551 (2015)



>> Structure - DMC(V) vs experiment

[a] J. Voigtsberger *et al.*, Nature Comm. 5, 5765 (2014)




❖ Summary

- Different scaling lengths were tested.
- Dimer, Borromean-trimer, all-bound trimer and HeHe-tango-trimer universal lines of scaled size-energy ratio were found.
- Clusters close to each other on universal line have similar structural properties.
- TTY+Ret potential trimer-distribution-predictions are in the best agreement with the experimental He-trimer data.
- For tetramers indication of different lines that depend on the number of bound subsystems of dimers was noticed.

❖ Acknowledgments

- Financial support:
 - ✓ Croatian Science Foundation (IP-2014-09-2452)
 - ❖ Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of Croatian Science Foundation.
 - ✓ MICINN-Spain (FIS2014-56257-C2-1-P)
- Computational support:



Thank you for attention! Questions?