

## Friedman (FLRW) universe

The *Friedman universe* is a cosmological model where the universe is assumed to be homogeneous, isotropic, and filled with *ideal fluid*.

The assumption of homogeneity and isotropicity of the universe is often referred to as the *cosmological principle*. Empirically, it seems to be justified on scales larger than 250 million light years. The isotropic and homogeneous model is sometimes called the *Standard Model* of present-day cosmology.

The *Friedman equation* is the Einstein equation applied to the Friedman universe as a whole. It describes the temporal evolution of a Friedman universe.

## Spaces with constant curvature

A homogeneous and isotropic universe is a *space with constant curvature*.

Let us first look at two-dimensional spaces of constant curvature: three-dimensional spheres (positive curvature), pseudo-spheres (negative curvature), and planes (zero curvature).

On a sphere the length element in the ordinary spherical coordinates is given as

$$dl^2 = a^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (1)$$

where  $a$  is the radius of the sphere.

Let us introduce the polar coordinates  $\{r, \phi\}$  on the sphere, with  $r$  measuring the distance to the north pole. The length of a circle around north pole,  $\theta = \text{const}$ , is equal  $2\pi a \sin \theta$ . Therefore if we want the circumference of the circle to be equal  $2\pi r$ , we need to define  $r = a \sin \theta$ .

The length element in the polar coordinates becomes

$$dl^2 = \frac{dr^2}{1 - \frac{r^2}{a^2}} + r^2 d\phi^2 . \quad (2)$$

On a pseudo-sphere, where the curvature is negative, the length element is correspondingly

$$dl^2 = \frac{dr^2}{1 + \frac{r^2}{a^2}} + r^2 d\phi^2 . \quad (3)$$

In angular coordinates  $r = a \sinh \theta$  the latter becomes

$$dl^2 = a^2 (d\theta^2 + \sinh^2 \theta d\phi^2) . \quad (4)$$

On a plane, where the curvature is zero, the length element is

$$dl^2 = dr^2 + r^2 d\phi^2 . \quad (5)$$

Analogously, a three-dimensional space with constant curvature can have one of the following three possible geometries:

- *flat* (zero curvature),

$$dl^2 = dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) = a^2 (d\chi^2 + \chi^2 (d\theta^2 + \sin^2 \theta d\phi^2)) , \quad (6)$$

where  $r = a\chi$ ,  $\chi \in [0, \infty]$ ;

- *closed* (positive curvature),

$$dl^2 = \frac{dr^2}{1 - \frac{r^2}{a^2}} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) = a^2 (d\chi^2 + \sin^2 \chi (d\theta^2 + \sin^2 \theta d\phi^2)) , \quad (7)$$

where  $r = a \sin \chi$ ,  $\chi \in [0, \pi]$ ;

- and *open* (negative curvature),

$$dl^2 = \frac{dr^2}{1 + \frac{r^2}{a^2}} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) = a^2 (d\chi^2 + \sinh^2 \chi (d\theta^2 + \sin^2 \theta d\phi^2)) , \quad (8)$$

where  $r = a \sinh \chi$ ,  $\chi \in [0, \infty[$ .

## Friedman equation

*Friedman metric*<sup>1</sup> is a (generic) synchronous metric in a homogeneous and isotropic universe,

$$ds^2 = dt^2 - a(t)^2 \left( d\chi^2 + \begin{bmatrix} \chi^2 \\ \sin^2 \chi \\ \sinh^2 \chi \end{bmatrix} (d\theta^2 + \sin^2 \theta d\phi^2) \right), \quad (9)$$

$a(t)$  is the time-dependent *scale factor* of the universe.

## Closed universe

In a closed Friedman universe the metric is

$$ds^2 = a^2 (d\eta^2 - d\chi^2 - \sin^2 \chi (d\theta^2 + \sin^2 \theta d\phi^2)), \quad (10)$$

where  $r = a \sin \chi$ , and  $\eta$  is the scaled time coordinate,

$$dt = a d\eta. \quad (11)$$

Now the following Maxima script

```
derivabbrev:true; /* a better notation for derivatives */
load(ctensor); /* load the package to deal with tensors */
ct_coords:[eta,chi,theta,phi]; /* our coordiantes: eta chi theta phi */
depends([a],[eta]); /* the scale, a, depends on eta */
lg:ident(4); /* set up a 4x4 identity matrix */
lg[1,1]: a^2; /* g_{\eta\eta} = a^2 */
lg[2,2]:-a^2; /* g_{\chi\chi} = -a^2 */
lg[3,3]:-a^2*sin(chi)^2; /* g_{\chi\chi} */
lg[4,4]:-a^2*sin(chi)^2*sin(theta)^2; /* g_{\phi\phi} */
cmetric(); /* compute the prerequisites for further calculations */
christof(mcs); /* print out the Christoffel symbols, mcs_{bca}=\Gamma^a_{bc} */
uricci(true); /* print out the elements of the Ricci tensor */
scurvature(); /* Ricci scalar curvature */
```

calculates the components of the Ricci tensor<sup>2</sup> and the Ricci scalar for this metric,

$$R_{\eta}^{\eta} = \frac{3}{a^4}(a'^2 - aa''), \quad R_{\chi}^{\chi} = R_{\theta}^{\theta} = R_{\phi}^{\phi} = -\frac{1}{a^4}(2a^2 + a'^2 + aa''), \quad R = -\frac{6}{a^3}(a + a''), \quad (12)$$

where prime denotes the  $\eta$ -derivative.

The stress-energy-momentum tensor for the perfect fluid is<sup>3</sup>

$$T_{ab} = (\epsilon + p)u_a u_b - p g_{ab}, \quad (13)$$

where  $\epsilon$  is the rest-energy density and  $p$  is the pressure. In synchronous Friedman coordinates the matter is at rest and the 4-velocity is  $u^b = \{\frac{1}{a}, 0, 0, 0\}$ .

<sup>1</sup>also referred to as Friedman-Lemaitre-Robertson-Walker in different combinations.

<sup>2</sup>The script also calculates the Christoffel symbols,  $\Gamma_{\eta\eta}^{\eta} = \Gamma_{\eta\chi}^{\chi} = \Gamma_{\eta\theta}^{\theta} = \Gamma_{\eta\phi}^{\phi} = \Gamma_{\chi\chi}^{\eta} = \frac{a'}{a}$ ,  $\Gamma_{\chi\theta}^{\theta} = \Gamma_{\chi\phi}^{\phi} = \cot \chi$ ,  $\Gamma_{\theta\theta}^{\eta} = \frac{a'}{a} \sin^2 \chi$ ,  $\Gamma_{\theta\theta}^{\chi} = -\cos \chi \sin \chi$ ,  $\Gamma_{\theta\phi}^{\phi} = \cot \theta$ ,  $\Gamma_{\phi\phi}^{\eta} = \frac{a'}{a} \sin^2 \chi \sin^2 \theta$ ,  $\Gamma_{\phi\phi}^{\chi} = -\cos \chi \sin \chi \sin^2 \theta$ ,  $\Gamma_{\phi\phi}^{\theta} = -\cos \theta \sin \theta$ .

<sup>3</sup>Interpreting the equations  $T^{ab}_{;b} = 0$  as conservation laws leads to the following interpretations of the components of the stress-energy-momentum tensor:  $T^{00}$  is energy density,  $T^{0\alpha}$  is momentum density,  $T^{\alpha\alpha}$  is pressure, and  $T^{\alpha\beta}$  where  $\alpha \neq \beta$  is the shear stress. For a perfect fluid at rest the shear stress is zero and the momentum is also zero. Thus in the frame where the element of the liquid is at rest,  $u^a = \{1, 0, 0, 0\}$ , the stress-energy-momentum tensor is diagonal with components  $\epsilon, p, p, p$  where  $\epsilon$  is the rest-energy and  $p$  is the pressure. Apparently, the covariant form must then be  $T^{ab} = (\epsilon + p)u^a u^b - p g^{ab}$ .

Correspondingly, the Einstein equation

$$R_b^a - \frac{1}{2}R\delta_b^a = \kappa T_b^a \quad (14)$$

then has the  $\frac{\eta}{\eta}$  component

$$\frac{3}{a^4}(a^2 + a'^2) = \kappa\epsilon, \quad (15)$$

and the three identical spatial components,

$$\frac{1}{a^4}(a^2 + 2aa'' - a'^2) = -\kappa p, \quad (16)$$

called the *Friedman equations* for a closed universe,

If the relation between  $\epsilon$  and  $p$ , called the *equation of state* of the matter, is known, the energy density  $\epsilon$  can be determined as function of  $a$  from the energy conservation equation. The latter must have the form

$$dE = -pdV, \quad (17)$$

where  $V$  is a volume element in the Friedman universe, and  $E = \epsilon V$  is the energy content of this volume. Since the volume is proportional to  $a^3$ , and both  $\epsilon$  and  $a$  in a Friedman universe can only depend on time, equation (17) can be rewritten as

$$(\epsilon a^3)' + p(a^3)' = 0. \quad (18)$$

It is easy to show, that equation (18) actually follows from the Friedman equations (15) and (16).

The energy conservation equation (17) can also be written as

$$\frac{3da}{a} = -\frac{d\epsilon}{\epsilon + p}. \quad (19)$$

When the dependence  $\epsilon(a)$  is found by integration of the energy conservation equation (19), the solution to the Friedman equation can be obtained as the integral

$$\eta = \pm \int \frac{da}{a\sqrt{\frac{1}{3}\kappa\epsilon a^2 - 1}}. \quad (20)$$

## Open universe

In an open Friedman universe the metric is

$$ds^2 = a^2 (d\eta^2 - d\chi^2 - \sinh^2 \chi (d\theta^2 + \sin^2 \theta d\phi^2)), \quad (21)$$

where  $r = a \sinh \chi$ , and  $ad\eta = dt$ . This metric can be obtained from the closed universe metric (10) by a formal substitution

$$\{a, \eta, \chi\} \rightarrow \{ia, i\eta, i\chi\}. \quad (22)$$

Therefore the Friedman equation for an open universe can be readily obtained from (15) by the substitution (22),

$$\frac{3}{a^4}(a'^2 - a^2) = \kappa\epsilon, \quad (23)$$

with the integral solution,

$$\eta = \pm \int \frac{da}{a\sqrt{\frac{1}{3}\kappa\epsilon a^2 + 1}}. \quad (24)$$

### Flat universe

In a flat Friedman universe the metric is

$$ds^2 = dt^2 - a(t)^2 (dx^2 + dy^2 + dz^2) , \quad (25)$$

and the Friedman equation is (see the exercise)

$$3 \frac{\dot{a}^2}{a^2} = \kappa \epsilon . \quad (26)$$

### Exercises

1. Our  $\eta$  Friedman equation is written for the  $\eta$ -derivative,  $a' \doteq da/d\eta$ . Rewrite it (as in the Wikipedia article) for the  $t$ -derivative  $\dot{a} \doteq da/dt$ .
2. Calculate manually the Ricci tensor in the Friedman coordinates for a closed universe.
3. Argue that the energy conservation equation,

$$(\epsilon a^3)' + p(a^3)' = 0 , \quad (27)$$

follows from the field equations

$$\frac{3}{a^4} (a^2 + a'^2) = \kappa \epsilon , \quad (28)$$

$$\frac{1}{a^4} (a^2 + 2aa'' - a'^2) = -\kappa p . \quad (29)$$

4. Consider a flat (Euclidean) isotropic universe with the metric

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2) .$$

and investigate its temporal development for matter and radiation dominated universes.

Hints:

- (a) Calculate the Christoffel symbols<sup>4</sup>;
  - (b) Calculate the Ricci tensor and the Ricci scalar<sup>5</sup>;
  - (c) Write down the  $t$  component of the Einstein equation with perfect fluid<sup>6</sup>;
  - (d) Write down the energy conservation equation<sup>7</sup>;
  - (e) Integrate the equations for a matter dominated universe ( $p = 0$ ,  $\epsilon = \mu$ )<sup>8</sup>;
  - (f) Integrate the equations for a radiation dominated universe ( $p = \epsilon/3$ )<sup>9</sup>;
5. Calculate the volumes of the closed and open universes.

---

4

$$\Gamma_{tx}^x = \Gamma_{ty}^y = \Gamma_{tz}^z = \frac{\dot{a}}{a} , \quad \Gamma_{xx}^t = \Gamma_{yy}^t = \Gamma_{zz}^t = a\dot{a} .$$

5

$$R_t^t = -3 \frac{\ddot{a}}{a} , \quad R_x^x = R_y^y = R_z^z = -\frac{\ddot{a}}{a} - 2 \frac{\dot{a}^2}{a^2} .$$

6

$$3 \frac{\dot{a}^2}{a^2} = \kappa \epsilon .$$

7

$$\frac{dV}{V} = -\frac{d\epsilon}{(\epsilon + p)} \Rightarrow 3 \ln(a) = -\int \frac{d\epsilon}{(\epsilon + p)} .$$

8

$$\mu a^3 = \text{const} , \quad a \propto t^{2/3} .$$

9

$$\epsilon a^4 = \text{const} , \quad a \propto t^{1/2} .$$